

University of Groningen

The role of cAMP-dependent protein kinase A in bile canalicular plasma membrane biogenesis in hepatocytes

Wojtal, Kacper Andrzej

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2007

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wojtal, K. A. (2007). *The role of cAMP-dependent protein kinase A in bile canalicular plasma membrane biogenesis in hepatocytes*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References

Abu-Absi, S. F., Hu, W. S., and Hansen, L. K. (2005) Dexamethasone effects on rat hepatocyte spheroid formation and function. *Tissue Eng.* 11, 415-426.

Aït Slimane, T., Trugnan, G., van IJzendoorn, S.C.D., and Hoekstra, D. (2003). Raft-mediated trafficking of apical resident proteins occurs in both direct and transcytotic pathways in polarized hepatic cells: role of distinct lipid microdomains. *Mol Biol Cell.* 14, 611-624.

Allan, V. J., Thompson, H. M., and McNiven, M. A. (2002) Motoring around the Golgi. *Nat Cell Biol.* 4, 236-242.

Alto, N.M., Soderling, S.H., Hoshi, N., Langeberg, L.K., Favos, R., Jennings, P.S., and Scott, J.D. (2003). Bioinformatic design of A-kinase anchoring protein-in silico: a potent and selective peptide antagonist of type II protein kinase A anchoring. *Proc Natl Acad Sci U S A.* 100, 4445-4450.

Alto, N., Carlisle Michel, J. J., Dodge, K. L., Langeberg, L. K., and Scott, J. D. (2002) Intracellular targeting of protein kinases and phosphatases. *Diabetes.* 51, 385-388.

Alto, N. M., Soderling, J., and Scott, J. D. (2002) Rab32 is an A-kinase anchoring protein and participates in mitochondrial dynamics. *J Cell Biol.* 158, 659-668.

Ameen, N. A., Marino, C., and Salas, P. J. (2003) cAMP-dependent exocytosis and vesicle traffic regulate CFTR and fluid transport in rat jejunum in vivo. *Am J Physiol Cell Physiol.* 284, 429-438.

Ameen, N. A., Martensson, B., Bourguignon, L., Marino, C., Isenberg, J., and McLaughlin, G. E. (1999) CFTR channel insertion to the apical surface in rat duodenal villus epithelial cells is upregulated by VIP in vivo. *J Cell Sci.* 112, 887-894.

Baba, T., Sakisaka, T., Mochida, S., and Takai, Y. (2005) PKA-catalyzed phosphorylation of tomosyn and its implication in Ca²⁺-dependent exocytosis of neurotransmitter. *J Cell Biol.* 170, 1113-1125.

Banales, J. M., Prieto, J., Medina, J. F. (2006) Cholangiocyte anion exchange and biliary bicarbonate excretion. *World J Gastroenterol.* 12, 3496-3511.

Bartles, J. R., Rao, M. S., Zhang, L. Q., Fayos, B. E., Nehme, C. L., and Reddy, J. K. (1991). Expression and compartmentalization of integral plasma membrane proteins by hepatocytes and their progenitors in the rat pancreas. *J Cell Sci.* 98, 45-54.

Barradeau, S., Imaizumi-Scherrer, T., Weiss, M, C., Faust, D. M. (2002) Intracellular targeting of the type-I alpha regulatory subunit of cAMP-dependent protein kinase. *Trends Cardiovasc Med.* 12, 235-241.

Bejarano, E., Cabrera, M., Vega, L., Hidalgo, J., and Velasco, A. (2006) Golgi structural stability and biogenesis depend on associated PKA activity. *J Cell Sci.* 119, 3764-3775.

Birkeli, K. A., Llorente, A., Torgersen, M. L., Keryer, G., Tasken, K., and Sandvig, K. (2003) Endosome-to-Golgi transport is regulated by protein kinase A type II alpha. *J Biol Chem.* 278, 1991-1997.

Boczan, J., Leenders, A. G., and Sheng, Z. H. (2004) Phosphorylation of syntaphilin by cAMP-dependent protein kinase modulates its interaction with syntaxin-1 and annuls its inhibitory effect on vesicle exocytosis. *J Biol Chem.* 279, 18911-18919.

Bos, J. L. (2003) Epac: a new cAMP target and new avenues in cAMP research. *Nat Rev Mol Cell Biol.* 4, 733-738.

Bouchard, J. F., Moore, S. W., Tritsch, N. X., Roux, P. P., Shekarabi, M., Barker, P. A., and Kennedy, T. E. (2004) Protein kinase A activation promotes plasma membrane insertion of DCC from an intracellular pool: A novel mechanism regulating commissural axon extension. *J Neurosci.* 24, 3040-3050.

Boucher, M. J., Laprise, P., and Rivard, N. (2005) Cyclic AMP-dependent protein kinase A negatively modulates adherens junction integrity and differentiation of intestinal epithelial cells. *J Cell Physiol.* 202, 178-190.

Birkeli, K. A., Llorente, A., Torgersen, M. L., Keryer, G., Tasken, K., and Sandvig, K. (2003) Endosome-to-Golgi transport is regulated by protein kinase A type II alpha. *J Biol Chem.* 278, 1991-1997.

Brignoni, M., Pignataro, O.P., Rodriguez, M.L., Alvarez, A., Vega-Salas, D.E., Rodriguez-Boulan, E., and Salas, P. J. (1995). Cyclic AMP modulates the rate of 'constitutive' exocytosis of apical membrane proteins in Madin-Darby canine kidney cells. *J Cell Sci.* 108, 1931-1943.

Brown, P.S., Wang, E., Aroeti, B., Chapin, S. J., Mostov, K. E., and Dunn, K. W. (2000) Definition of distinct compartments in polarized Madin-Darby canine kidney (MDCK) cells for membrane-volume sorting, polarized sorting and apical recycling. *Traffic.* 1, 124-140.

Buendia, B., Bre, M. H., Griffiths, G., and Karsenti, E. (1990) Cytoskeletal control of centrioles movement during the establishment of polarity in Madin-Darby canine kidney cells. *J Cell Biol.* 110, 1123-1135.

Burakov, A., Nadezhdina, E., Slepchenko, B., and Rodionov, V. (2003) Centrosome positioning in interphase cells. *J Cell Biol.* 162, 963-969.

Burgos, P.V., Klattenhoff, C., de la Fuente, E., Rigotti, A., and Gonzalez, A. (2004). Cholesterol depletion induces PKA-mediated basolateral-to-apical

transcytosis of the scavenger receptor class B type I in MDCK cells. *Proc Natl Acad Sci U S A.* 101, 3845-3850.

Butterworth, M.B., Helman, S.I., and Els, W.J. (2001). cAMP-sensitive endocytic trafficking in A6 epithelia. *Am J Physiol Cell Physiol.* 280, 752-762.

Buterworth, M.B., Edinger, R.S., Johnson, J.P., and Frizzel, R. A. (2005). Acute ENaC stimulation by cAMP in a kidney cell line is mediated by exocytic insertion from a recycling channel pool. *J Gen Physiol.* 125, 81-101.

Cabrera, M., Muniz, M., Hidalgo, J., Vega, L., Martin, M. E., and Velasco, A. (2003) The retrieval function of the KDEL receptor requires PKA phosphorylation of its C-terminus. *Mol Biol Cell.* 14, 4114-4125.

Cantz, T., Nies, A.T., Brom, M., Hofmann, A.F., and Keppler, D. (2000). MRP2, a human conjugate export pump, is present and transports fluo 3 into apical vacuoles of Hep G2 cells. *Am J Physiol Gastrointest Liver Physiol.* 278, 522-531.

Carlson, C. R., Lygren, B., Berge, T., Hoshi, N., Wong, W., Tasken, K., and Scott, J. D. (2006) Delineation of type I protein kinase A-selective signaling events using an RI anchoring disruptor. *J Biol Chem.* 281, 21535-21545.

Carlson, C. R., Witczak, O., Vossebein, L., Labbé, J. C., Skálhegg, B. S., Keryer, G., Herberg, F. W., Collas, P., and Taskén, K. (2001) CDK1-mediated phosphorylation of the RII α regulatory subunit of PKA works as a molecular switch that promotes dissociation of RII α from centrosomes at mitosis. *J. Cell Sci.* 114, 3243-3254.

Carmosino, M., Procino, G., Tamma, G., Mannucci, R., Svelto, M., Valenti, G. (2006) Trafficking and phosphorylation dynamics of AQP4 in histamine-treated human gastric cells. *Biol Cell.* [Epub ahead of print]

Carr, D. W., Stofko-Hahn, R. E., Fraser, I. D., Bishop, S. M., Acott, T. S., Brennan, R. G., and Scott, J. D. (1991) Interaction of the regulatory subunit (RII) of cAMP-dependent protein kinase with RII-anchoring proteins occurs through an amphipathic helix binding motif. *J Biol Chem.* 266, 14188-14192.

Casanova, J. E., Wang, X., Kumar, R., Bhartur, S. G., Navarre, J., Woodrum, J. E., Altschuler, Y., Ray, G. S., and Goldenring, J. R. (1999) Association of Rab25 and Rab11a with the apical recycling system of polarized Madin-Darby canine kidney cells. *Mol Biol Cell.* 10, 47-61.

Chang, S. Y., Di, A., Naren, A. P., Palfrey, H. C., Kirk, K. L., Nelson, D. J. (2002) Mechanisms of CFTR regulation by syntaxin 1A and PKA. *J Cell Sci.* 115, 783-791.

Chappe, V., Irvine, T., Liao, J., Evagelidis, A., and Hanrahan, J. W. (2005) Phosphorylation of CFTR by PKA promotes binding of the regulatory domain. *EMBO J.* 24, 2730-2740.

Chen, X., Dai, J. C., Orellana, S. A., and Greenfield, E. M. (2005) Endogenous protein kinase inhibitor gamma terminates immediate-early gene expression induced by cAMP-dependent protein kinase (PKA) signaling: termination depends on PKA inactivation rather than PKA export from the nucleus. *J Biol Chem.* 280, 2700-2707.

Chin, K. V., Yang, W. L., Ravatn, R., Kita, T., Reitman, E., Vettori, D., Cvijic, M. E., Shin, M., and Iacono, L. (2002) Reinventing the wheel of cyclic AMP: novel mechanisms of cAMP signaling. *Ann N Y Acad Sci.* 968, 49-64.

Christensen, A. E., Selheim, F., de Rooij, J., Dremier, S., Schwede, F., Dao, K. K., Martinez, A., Maenhaut, C., Bos, J. L., Genieser, H. G., and Døskeland, S. O. (2003) cAMP analog mapping of Epac1 and cAMP kinase. Discriminating analogs demonstrate that Epac and cAMP kinase act synergistically to promote PC-12 cell neurite extension. *J Biol Chem.* 278, 35394-35402.

Cohen, D., Brennwald, P. J., Rodriguez-Boulán, E., and Musch, A. (2004) Mammalian PAR-1 determines epithelial lumen polarity by organizing the microtubule cytoskeleton. *J Cell Biol.* 164, 717-727.

Constantinescu, A., Diamond, I., Gordon, A. S. (1999) Ethanol-induced translocation of cAMP-dependent protein kinase to the nucleus. Mechanism and functional consequences. *J Biol Chem.* 274, 26985-26991.

Cowan, C. R., and Hyman, A. A. (2004) Centrosomes direct cell polarity independently of microtubule assembly in *C. elegans* embryos. *Nature.* 431, 92-96.

Colledge, M., and Scott, J.D. (1999). AKAPs: from structure to function. *Trends Cell Biol.* 9, 216-21.

Conti, M. (2000) Phosphodiesterases and cyclic nucleotide signaling in endocrine cells. *Mol Endocrinol.* 14, 1317-1327.

Diviani, D., and Scott, J. D. (2001) AKAP signaling complexes at the cytoskeleton. *J Cell Sci.* 114, 1431-1437.

D'Souza, T., Agarwal, R., and Morin, P. J. (2005) Phosphorylation of claudin-3 at threonine 192 by cAMP-dependent protein kinase regulates tight junction barrier function in ovarian cancer cells. *J Biol Chem.* 280, 26233-26240.

D'Souza-Schorey, C., Chavrier, P. (2006) ARF proteins: roles in membrane traffic and beyond. *Nat Rev Mol Cell Biol.* 7, 347-358.

de Anda, F.C., Pollarolo, G., Da Silva, J. S., Camoletto, P., G., Feiguin, F., and Dotti, C. G. (2005) Centrosome localization determines neuronal polarity. *Nature*. 436, 704-708.

de Matteis MA, Luna A, Di Tullio G, Corda D, Kok JW, Luini A, Egea G. (1999). PDMP blocks the BFA-induced ADP-ribosylation of BARS-50 in isolated Golgi membranes. *FEBS Lett*. 459, 310-312.

de Rooij, J., Zwartkruis, F. J., Verheijen, M. H., Cool, R. H., Nijman, S. M., Wittinghofer, A., Bos, J. L. (1998) Epac is a Rap1 guanine-nucleotide-exchange factor directly activated by cyclic AMP. *Nature*. 396, 474-477.

Dijkhuis, A.J., Douwes, J., Kamps, W., Sietsma, H., and Kok, J.W. (2003). Differential expression of sphingolipids in P-glycoprotein or multidrug resistance-related protein 1 expressing human neuroblastoma cell lines. *FEBS Lett*. 548, 28-32.

Diviani, D. and Scott, J. D. (2001) AKAP signaling complexes at the cytoskeleton *J.Cell Sci*. 114, 1431-1437.

Diviani, D., Langeberg, L. K, Doxsey, S. and Scott, J. D. (2000) Pericentrin anchors protein kinase A at the centrosomes through a newly identified RII-binding domain *Curr. Biol*. 10, 417-420.

Dostmann, W. R., Taylor, S. S., Genieser, H. G., Jastorff, B., Doskeland, S. O., and Ogreid, D. (1990) Probing the cyclic nucleotide binding sites of cAMP-dependent protein kinases I and II with analogs of adenosine 3',5'-cyclic phosphorothioates. *J Biol Chem*. 265, 10484-1091.

Doxsey, S. J., Stein, P., Evans, L., Calarco, P. D., and Kirschner, M. (1994) Pericentrin, a highly conserved centrosome protein involved in microtubule organization. *Cell*. 76, 639-650.

Dylewski, D. P., and Keenan, T. W. (1984) Centrioles in the mammary epithelium of the rat. *J Cell Sci*. 72, 185-193.

Edwards, A.S., and Scott, J.D. (2000). A-kinase anchoring proteins: protein kinase A and beyond. *Curr Opin Cell Biol*. 12, 217-221.

Ekanger, R., Vintermyr, O. K., Houge, G., Sand, T. E., Scott, J. D., Krebs, E. G., Eikhom, T. S., Christoffersen, T., Ogreid, D., and Doskeland, S. O. (1989) The expression of cAMP-dependent protein kinase subunits is differentially regulated during liver regeneration. *J Biol Chem*. 264, 4374-4382.

Evans, G. J., and Morgan, A. (2003) Regulation of the exocytotic machinery by cAMP-dependent protein kinase: implications for presynaptic plasticity. *Biochem Soc Trans*. 31, 824-827.

Falasca, L., Tisone, G., Palmieri, G., Anselmo, A., Di Paolo, D., Baiocchi, L., Torri, E., Orlando, G., Casciani, C. U., and Angelico, M. (2001) Protective role of tauroursodeoxycholate during harvesting and cold storage of human liver: a pilot study in transplant recipients. *Transplantation*. 71, 1268-1276.

Fath, K. R., Trimbur, G. M., and Burgess, D. R. (1994) Molecular motors are differentially distributed on Golgi membranes from polarized epithelial cells. *J Cell Biol*. 126, 661-675.

Feliciello, A., Gallo, A., Mele, E., Porcellini, A., Troncone, G., Garbi, C., Gottesman, M. E., and Avvedimento, E. V. (2000) The localization and activity of cAMP-dependent protein kinase affect cell cycle progression in thyroid cells. *J Biol Chem*. 275, 303-311.

Garcia, F., Kierbel, A.M., Larocca, M.C., Gradilone, S.A., Splinter, P., LaRusso, N.F., and Marinelli, R.A. (2001). The water channel aquaporin-8 is mainly intracellular in rat hepatocytes, and its plasma membrane insertion is stimulated by cyclic AMP. *J Biol Chem*. 276, 12147-12152.

Genieser, H. G., Winkler, E., Butt, E., Zorn, M., Schulz, S., Iwitzki, F., Stormann, R., Jastorff, B., Doskeland, S. O., OGREID, D., et al. (1992) Derivatives of 1-beta-D-ribofuranosylbenzimidazole 3',5'-phosphate that mimic the actions of adenosine 3',5'-phosphate (cAMP) and guanosine 3',5'-phosphate (cGMP). *Carbohydr Res*. 234, 217-235.

Golub, T., and Caroni, P. (2005) PI(4,5)P2-dependent microdomain assemblies capture microtubules to promote and control leading edge motility. *J Cell Biol*. 169, 151-165.

Gorski, J. A., Gomez, L. L., Scott, J. D., and Dell'Acqua, M. L. (2005) Association of an A-kinase-anchoring protein signaling scaffold with cadherin adhesion molecules in neurons and epithelial cells. *Mol Biol Cell*. 16, 3574-3590.

Gradilone, S. A., Garcia, F., Huebert, R. C., Tietz, P. S., Larocca, M. C., Kierbel, A., Carreras, F. I., Larusso, N. F., and Marinelli, R. A. (2003). Glucagon induces the plasma membrane insertion of functional aquaporin-8 water channels in isolated rat hepatocytes. *Hepatology*. 37, 1435-1441.

Grindstaff, K. K., Bacallao, R. L., and Nelson, W. J. (1998) Apiconuclear organization of microtubules does not specify protein delivery from the trans-Golgi network to different membrane domains in polarized epithelial cells. *Mol Biol Cell*. 9, 685-699.

Hansen, S. H., Oleson, A., and Casanova, J. E. (1995). Wortmannin, an inhibitor of phosphoinositide 3-kinase, inhibits transcytosis in polarized epithelial cells. *J Biol Chem*. 270, 28425-28432.

Hansen, S. H., and Casanova, J. E. (1994) Gs alpha stimulates transcytosis and apical secretion in MDCK cells through cAMP and protein kinase A. *J Cell Biol.* 126, 677-687.

Howard, M., Jiang, X., Stoltz, D. B., Hill, W. G., Johnson, J. A., Watkins, S. C., Frizzel, R. A., Bruton, C. M., Robbins, P. D., and Weisz, O. A. (2000). Forskolin-induced apical membrane insertion of virally expressed, epitope-tagged CFTR in polarized MDCK cells. *Am J Physiol Cell Physiol.* 279, 375-382.

Howard, M., Jilling, T., DuVall, M., Frizzell, R. A. (1996) cAMP-regulated trafficking of epitope-tagged CFTR. *Kidney Int.* 49, 1642-1648.

Henn, V., Edemir, B., Stefan, E., Wiesner, B., Lorenz, D., Theilig, F., Schmitt, R., Vossebein, L., Tamma, G., Beyermann, M., Krause, E., Herberg, F.W., Valenti, G., Bachmann, S., Rosenthal, W., and Klusmann, E. (2004). Identification of a novel A-kinase anchoring protein 18 isoform and evidence for its role in the vasopressin-induced aquaporin-2 shuttle in renal principal cells. *J Biol Chem.* 279, 26654-26665.

Herberg, F. W., Maleszka, A., Eide, T., Vossebein, L., and Tasken, K. (2000) Analysis of A-kinase anchoring protein (AKAP) interaction with protein kinase A (PKA) regulatory subunits: PKA isoform specificity in AKAP binding. *J Mol Biol.* 298, 329-339.

Herrema, H., Czajkowska, D., Theard, D., van der Wouden, J. M., Kalicharan, D., Zolghadr, B., Hoekstra, D., and van Ijzendoorn, S. C. (2006) Rho Kinase, Myosin-II, and p42/44 MAPK Control Extracellular Matrix-mediated Apical Bile Canalicular Lumen Morphogenesis in HepG2 Cells. *Mol Biol Cell.* 17, 3291-3303.

Higginbotham, H., Tanaka, T., Brinkman, B. C., and Gleeson, J. G. (2006) GSK3beta and PKCzeta function in centrosome localization and process stabilization during Slit-mediated neuronal repolarization. *Mol Cell Neurosci.* 32, 118-132.

Hinrichs, J. W., Klappe, K., Hummel, I., and Kok, J. W. (2004). ATP-binding cassette transporters are enriched in non-caveolar detergent-insoluble glycosphingolipid-enriched membrane domains (DIGs) in human multidrug-resistant cancer cells. *J Biol Chem.* 279, 5734-5738.

Hobdy-Henderson, K. C., Hales, C. M., Lapierre, L. A., Cheney, R. E., and Goldenring, J. R. (2003) Dynamics of the apical plasma membrane recycling system during cell division. *Traffic.* 4, 681-693.

Hoekstra, D., Tyteca, D., and van Ijzendoorn, S. C. (2004) The subapical compartment: a traffic center in membrane polarity development. *J Cell Sci.* 117, 2183-2192.

Howe, A. K., Baldor, L. C., and Hogan, B. P. (2005) Spatial regulation of the cAMP-dependent protein kinase during chemotactic cell migration. *Proc Natl Acad Sci U S A.* 102, 14320-14325.

Huang, L. J., Wang, L., Ma, Y., Durick, K., Perkins, G., Deerinck, T. J., Ellisman, M. H., and Taylor, S. S. (1999) NH₂-Terminal targeting motifs direct dual specificity A-kinase-anchoring protein 1 (D-AKAP1) to either mitochondria or endoplasmic reticulum. *J Cell Biol.* 145, 951-959.

Huang, P., Gilmore, E., Kultgen, P., Barnes, P., Milgram, S., and Stutts, M. J. (2004) Local regulation of cystic fibrosis transmembrane regulator and epithelial sodium channel in airway epithelium. *Proc Am Thorac Soc.* 1, 33-37.

Huang, P., Trotter, K., Boucher, R. C., Milgram, S. L., and Stutts, M. J. (2000) PKA holoenzyme is functionally coupled to CFTR by AKAPs. *Am J Physiol Cell Physiol.* 278, 417-422.

Hundsruker, C., Rosenthal, W., and Klussmann, E. (2006) Peptides for disruption of PKA anchoring. *Biochem Soc Trans.* 34, 472-473.

Hundsruker, C., Krause, G., Beyermann, M., Prinz, A., Zimmermann, B., Diekmann, O., Lorenz, D., Stefan, E., Nedvetsky, P., Dathe, M., Christian, F., McSorley, T., Krause, E., McConnachie, G., Herberg, F. W., Scott, J. D., Rosenthal, W., and Klussmann, E. (2006) High-affinity AKAP7delta-protein kinase A interaction yields novel protein kinase A-anchoring disruptor peptides. *Biochem J.* 396, 297-306.

Ihrke, G., Neufeld, E. B., Meads, T., Shanks, M. R., Cassio, D., Laurent, M., Schroer, T. A., Pagano, R. E., Hubbard, A. L. (1993) WIF-B cells: an in vitro model for studies of hepatocyte polarity. *J Cell Biol.* 123, 1761-1775.

Jiang, S., Rhee, S. W., Gleeson, P. A., and Storrie, B. (2006) Capacity of the Golgi apparatus for cargo transport prior to complete assembly. *Mol Biol Cell.* 17, 4105-4117.

Jo, I., Ward, D.T., Baum, M.A., Scott, J.D., Coghlan, V.M., Hammond, T.G., and Harris, H.W. (2001). AQP2 is a substrate for endogenous PP2B activity within an inner medullary AKAP-signaling complex. *Am J Physiol Renal Physiol.* 281, 958-965.

Kagawa, T., Varticovski, L., Sai, Y., and Arias, I.M. (2002). Mechanism by which cAMP activates PI3-kinase and increases bile acid secretion in WIF-B9 cells. *Am J Physiol Cell Physiol.* 283, 1655-1666.

Kamiya, A., Kinoshita, T., and Miyajima, A. (2001) Oncostatin M and hepatocyte growth factor induce hepatic maturation via distinct signaling pathways. *FEBS Lett.* 492, 90-94.

Kamiya, A., Kinoshita, T., Ito, Y., Matsui, T., Morikawa, Y., Senba, E., Nakashima, K., Taga, T., Yoshida, K., Kishimoto, T., and Miyajima, A.

(1999) Fetal liver development requires a paracrine action of oncostatin M through the gp130 signal transducer. *EMBO J.* 18, 2127-2136.

Kang, G., Chepurny, O. G., Malester, B., Rindler, M. J., Rehmann, H., Bos, J. L., Schwede, F., Coetzee, W. A., and Holz, G. G. (2006) cAMP sensor Epac as a determinant of ATP-sensitive potassium channel activity in human pancreatic beta cells and rat INS-1 cells. *J Physiol.* 573, 595-609.

Karvar, S., Yao, X., Duman, J. G., Hybiske, K., Liu, Y., and Forte, J.G. (2002). Intracellular distribution and functional importance of vesicle-associated membrane protein 2 in gastric parietal cells. *Gastroenterology.* 123, 281-290.

Kashina, A. S., Semenova, I.V., Ivanov, P. A., Potekhina, E. S., Zaliapin, I., Rodionov, V., I. (2004) Protein kinase A, which regulates intracellular transport, forms complexes with molecular motors on organelles. *Curr Biol.* 14, 1877-1881.

Kato, A., Gores, G. J., and LaRusso, N. F. (1992) Secretin stimulates exocytosis in isolated bile duct epithelial cells by a cyclic AMP-mediated mechanism. *J Biol Chem.* 267, 15523-15529.

Katsura, T., Gustafson, C. E., Ausiello, D. A., and Brown, D. (1997) Protein kinase A phosphorylation is involved in regulated exocytosis of aquaporin-2 in transfected LLC-PK1 cells. *Am J Physiol.* 272, 817-822.

Kelly, J. H., and Darlington, G. J. (1989) Modulation of the liver specific phenotype in the human hepatoblastoma line Hep G2. *In Vitro Cell Dev Biol.* 25, 217-222.

Keryer, G., Skalhegg, B. S., Landmark, B. F., Hansson, V., Jahnsen, T., and Tasken, K. (1999) Differential localization of protein kinase A type II isozymes in the Golgi-centrosomal area. *Exp Cell Res.* 249, 131-146.

Keryer, G., Alsat, E., Tasken, K., and Evain-Brion, D. (1998) Cyclic AMP-dependent protein kinases and human trophoblast cell differentiation in vitro. *J Cell Sci.* 111, 995-1004.

Keryer, G., Luo, Z., Cavadore, J. C., Erlichman, J., and Bornens, M. (1993) Phosphorylation of the regulatory subunit of type II beta cAMP-dependent protein kinase by cyclin B/p34cdc2 kinase impairs its binding to microtubule-associated protein 2. *Proc Natl Acad Sci U S A.* 90, 5418-5422.

Kinoshita, T., and Miyajima, A. (2002). Cytokine regulation of liver development. *Biochim Biophys Acta.* 1592, 303-312.

Kipp, H., Pichetshote, N., and Arias, I. M. (2001). Transporters on demand: intrahepatic pools of canalicular ATP binding cassette transporters in rat liver. *J Biol Chem.* 276, 7218-7224.

Kipp, H., and Arias, I.M. (2000). Newly synthesized canalicular ABC transporters are directly targeted from the Golgi to the hepatocyte apical domain in rat liver. *J Biol Chem.* 275, 15917-15925.

Kiermayer, S., Biondi, R. M., Imig, J., Plotz, G., Haupenthal, J., Zeuzem, S., and Piiper, A. (2005) Epac activation converts cAMP from a proliferative into a differentiation signal in PC12 cells. *Mol Biol Cell.* 16, 5639-5648.

Klappe, K., Hinrichs, J. W., Kroesen, B. J., Sietsma, H., and Kok, J. W. (2004). MRP1 and glucosylceramide are coordinately over expressed and enriched in rafts during multidrug resistance acquisition in colon cancer cells. *Int J Cancer.* 110, 511-522.

Klausner, R. D., Donaldson, J. G., and Lippincott-Schwartz, J. (1992) Brefeldin A: insights into the control of membrane traffic and organelle structure. *J Cell Biol.* 116, 1071-1080.

Kleizen, B., Braakman, I., and de Jonge, H. R. (2000) Regulated trafficking of the CFTR chloride channel. *Eur J Cell Biol.* 79, 544-556.

Klussmann, E., and Rosenthal, W. (2001) Role and identification of protein kinase A anchoring proteins in vasopressin-mediated aquaporin-2 translocation. *Kidney Int.* 60, 446-449.

Klussmann, E., Maric, K., Wiesner, B., Bevermann, M., and Rosenthal W. (1999). Protein kinase A anchoring proteins are required for vasopressin-mediated translocation of aquaporin-2 into cell membranes of renal principal cells. *J Biol Chem.* 274, 4934-4938.

Knepper, M.A., and Inoue, T. (1997). Regulation of aquaporin-2 water channel trafficking by vasopressin. *Curr Opin Cell Biol.* 9, 560-564.

Kok JW, Babia T, Filipeanu CM, Nelemans A, Egea G, Hoekstra D. (1998). PDMP blocks brefeldin A-induced retrograde membrane transport from golgi to ER: evidence for involvement of calcium homeostasis and dissociation from sphingolipid metabolism. *J Cell Biol.* 142, 25-38.

Kraemer, J., Schmitz, F., and Drenckhahn, D. (1999) Cytoplasmic dynein and dynactin as likely candidates for microtubule-dependent apical targeting of pancreatic zymogen granules. *Eur J Cell Biol.* 78, 265-277.

Kubitz, R., Huth, C., Schmitt, M., Horbach, A., Kullak-Ublick, G., and Haussinger, D. (2001). Protein kinase C-dependent distribution of the multidrug resistance protein 2 from the canalicular to the basolateral membrane in human HepG2 cells. *Hepatology.* 34, 340-350.

Kopperud, R., Krakstad, C., Selheim, F., Doskeland, S. O. (2003) cAMP effector mechanisms. Novel twists for an 'old' signaling system. *FEBS Lett.* 546, 121-126.

Lacana, E., Maceyka, M., Milstien, S., and Spiegel, S. (2002) Cloning and characterization of a protein kinase A anchoring protein (AKAP)-related protein that interacts with and regulates sphingosine kinase 1 activity. *J Biol Chem.* 277, 32947-32953.

Lane, J. D., Lucocq, J., Pryde, J., Barr, F. A., Woodman, P. G., Allan, V. J., and Lowe, M. (2002) Caspase-mediated cleavage of the stacking protein GRASP65 is required for Golgi fragmentation during apoptosis. *J Cell Biol.* 156, 495-509.

Lande, M. B., Jo, I., Zeidel, M. L., Somers, M., Harris, and H. W. Jr. (1996) Phosphorylation of aquaporin-2 does not alter the membrane water permeability of rat papillary water channel-containing vesicles. *J Biol Chem.* 271, 5552-5557.

Lange, B. M. (2002) Integration of the centrosome in cell cycle control, stress response and signal transduction pathways. *Curr Opin Cell Biol.* 14, 35-43.

Lapierre, L. A., Kumar, R., Hales, C. M., Navarre, J., Bhartur, S. G., Burnette, J. O., Provance, D. W. Jr., Mercer, J. A., Bahler, M., and Goldenring, J. R. (2001) Myosin vb is associated with plasma membrane recycling systems. *Mol Biol Cell.* 12, 1843-1857.

Larocca, M. C., Shanks, R. A., Tian, L., Nelson, D. L., Stewart, D. M., and Goldenring, J. R. (2004) AKAP350 interaction with cdc42 interacting protein 4 at the Golgi apparatus. *Mol Biol Cell.* 15, 2771-2781.

Larocca, M. C., Jin, M., and Goldenring, J. R. (2006) AKAP350 modulates microtubule dynamics. *Eur J Cell Biol.* 85, 611-619.

Lavie, Y., Cao, H., Bursten, S.L., Giuliano, A.E., and Cabot, M.C. (1996). Accumulation of glucosylceramides in multidrug-resistant cancer cells. *J Biol Chem.* 271, 19530-19536.

Lavie, Y., and Liscovitch, M. (2000). Changes in lipid and protein constituents of rafts and caveolae in multidrug resistant cancer cells and their functional consequences. *Glycoconj J.* 17, 253-259.

LeCluyse, E. L., Audus, K. L., and Hochman, J. H. (1994) Formation of extensive canalicular networks by rat hepatocytes cultured in collagen-sandwich configuration. *Am J Physiol.* 266, 1764-1774.

Lester, L. B., Langeberg, L. K., and Scott, J. D. (1997) Anchoring of protein kinase A facilitates hormone-mediated insulin secretion. *Proc Natl Acad Sci U S A.* 94, 14942-14947.

Leung, S. M., Ruiz, W. G., and Apodaca, G. (2000) Sorting of membrane and fluid at the apical pole of polarized Madin-Darby canine kidney cells. *Mol Biol Cell.* 11, 2131-2150.

Li, H., Adamik, R., Pacheco-Rodriguez, G., Moss, J., and Vaughan, M. (2003) Protein kinase A-anchoring (AKAP) domains in brefeldin A-inhibited guanine nucleotide-exchange protein 2 (BIG2). *Proc Natl Acad Sci U S A.* 100, 1627-1632.

Llorente, A., van Deurs, B., and Sandvig, K. (1998) Transport of apically but not basolaterally internalized ricin to the Golgi apparatus is stimulated by 8-Br-cAMP in MDCK cells. *FEBS Lett.* 431, 200-204.

Lukacs, G. L., Segal, G., Kartner, N., Grinstein, S., and Zhang, F. (1997) Constitutive internalization of cystic fibrosis transmembrane conductance regulator occurs via clathrin-dependent endocytosis and is regulated by protein phosphorylation. *Biochem J.* 328, 353-361.

Ma, Y., Pitson, S., Hercus, T., Murphy, J., Lopez, A., and Woodcock, J. (2005) Sphingosine activates protein kinase A type II by a novel cAMP-independent mechanism. *J Biol Chem.* 280, 26011-26017.

Maier, O., and Hoekstra, D. (2003). Trans-Golgi network and subapical compartment of HepG2 cells display different properties in sorting and exiting of sphingolipids. *J Biol Chem.* 278, 164-173.

Martin, M. E., Hidalgo, J., Rosa, J. L., Crottet, P., and Velasco, A. (2000) Effect of protein kinase A activity on the association of ADP-ribosylation factor 1 to golgi membranes. *J Biol Chem.* 275, 19050-19059.

Martin, M. E., Hidalgo, J., Vega, F. M., and Velasco, A. (1999) Trimeric G proteins modulate the dynamic interaction of PKAII with the Golgi complex. *J Cell Sci.* 112, 3869-3878.

Mavrakis, M., Lippincott-Schwartz, J., Stratakis, C. A., and Bossis, I. (2006) Depletion of type IA regulatory subunit (RI α) of protein kinase A (PKA) in mammalian cells and tissues activates mTOR and causes autophagic deficiency. *Hum Mol Genet.* 15, 2962-2971.

McSorley T, Stefan E, Henn V, Wiesner B, Baillie GS, Houslay MD, Rosenthal W, Klussmann E. (2006) Spatial organisation of AKAP18 and PDE4 isoforms in renal collecting duct principal cells. *Eur J Cell Biol.* 85, 673-678.

Meads, T. and Schroer, T. A. (1995) Polarity and nucleation of microtubules in polarized epithelial cells. *Cell Motil. Cytoskel.* 32, 273-288.

Michel, J.J., and Scott, J.D. (2002). AKAP mediated signal transduction. *Annu Rev Pharmacol Toxicol.* 42, 235-257.

Misra, S., Varticovski, L., and Arias, I. M. (2003) Mechanisms by which cAMP increases bile acid secretion in rat liver and canalicular membrane vesicles. *Am J Physiol Gastrointest Liver Physiol.* 285, 316-324.

Misra, S., Ujhazy, P., Gatmaitan, Z., Varticovski, L., and Arias, I. M. (1998) The role of phosphoinositide 3-kinase in taurocholate-induced trafficking of ATP-dependent canalicular transporters in rat liver. *J Biol Chem.* 273, 26638-26644.

Miyajima, A., Kinoshita, T., Tanaka, M., Kamiya, A., Mukouyama, Y., and Hara, T. (2000). Role of Oncostatin M in hematopoiesis and liver development. *Cytokine Growth Factor Rev.* 11, 177-183.

Mogelsvang, S., Marsh, B. J., Ladinsky, M. S., Howell, K. E. (2004) Predicting function from structure: 3D structure studies of the mammalian Golgi complex. *Traffic.* 5, 338-345.

Mordasini, D., Bustamante, M., Rousselot, M., Martin, P. Y., Hasler, U., and Feraille, E. (2005) Stimulation of Na⁺ transport by AVP is independent of PKA phosphorylation of the Na-K-ATPase in collecting duct principal cells. *Am J Physiol Renal Physiol.* 289, 1031-1039.

Mundell, S. J., Pula, G., More, J. C., Jane, D. E., Roberts, P. J., and Kelly, E. (2004) Activation of cyclic AMP-dependent protein kinase inhibits the desensitization and internalization of metabotropic glutamate receptors 1a and 1b. *Mol Pharmacol.* 65, 1507-1516.

Muniz, M., Martin, M. E., Hidalgo, J., and Velasco, A. (1997) Protein kinase A activity is required for the budding of constitutive transport vesicles from the trans-Golgi network. *Proc Natl Acad Sci U S A.* 94, 14461-14466.

Muniz, M., Alonso, M., Hidalgo, J., Velasco, A. (1996) A regulatory role for cAMP-dependent protein kinase in protein traffic along the exocytic route. *J Biol Chem.* 271, 30935-30941.

Musat, A. I., Sattler, C. A., Sattler, G. L., and Pitot, H. C. (1993) Reestablishment of cell polarity of rat hepatocytes in primary culture. *Hepatology.* 18, 198-205.

Nejsum, L. N., Zelenina, M., Aperia, A., Frokiaer, J., and Nielsen, S. (2005) Bidirectional regulation of AQP2 trafficking and recycling: involvement of AQP2-S256 phosphorylation. *Am J Physiol Renal Physiol.* 288, 930-938.

Noda, Y., Okada, Y., Saito, N., Setou, M., Xu, Y., Zhang, Z., and Hirokawa, N. (2001) KIFC3, a microtubule minus end-directed motor for the apical transport of annexin XIIIb-associated Triton-insoluble membranes. *J Cell Biol.* 155, 77-88.

O'Connor, K. L., Mercurio, A. M. (2001) Protein kinase A regulates Rac and is required for the growth factor-stimulated migration of carcinoma cells. *J Biol Chem.* 276, 47895-47900.

Ogreid, D., Dostmann, W., Genieser, H. G., Niemann, P., Doskeland, S. O., and Jastorff, B. (1994) (Rp)- and (Sp)-8-piperidino-adenosine 3',5'-

(cyclic)thiophosphates discriminate completely between site A and B of the regulatory subunits of cAMP-dependent protein kinase type I and II. *Eur J Biochem.* 221, 1089-1094.

Ogreid, D., Ekanger, R., Suva, R. H., Miller, J. P., and Doskeland, S. O. (1989) Comparison of the two classes of binding sites (A and B) of type I and type II cyclic-AMP-dependent protein kinases by using cyclic nucleotide analogs. *Eur J Biochem.* 181, 19-31.

Peng, X., Grosse, B., Le Tiec, B., Nicolas, V., Delagebeaudeuf, C., Bedda, T., Decaens, C., and Cassio, D. (2006) How to induce non-polarized cells of hepatic origin to express typical hepatocyte polarity: generation of new highly polarized cell models with developed and functional bile canaliculi. *Cell Tissue Res.* 323, 233-243.

Perez, L. M., Milkiewicz, P., Ahmed-Choudhury, J., Elias, E., Ochoa, J. E., Sanchez Pozzi, E.J., Coleman, R., and Roma, M. G. (2006) Oxidative stress induces actin-cytoskeletal and tight-junctional alterations in hepatocytes by a Ca^{2+} -dependent, PKC-mediated mechanism: protective effect of PKA. *Free Radic Biol Med.* 40, 2005-2017.

Perez, L. M., Milkiewicz, P., Elias, E., Coleman, R., Sanchez Pozzi, E. J., and Roma, M. G. (2006) Oxidative stress induces internalization of the bile salt export pump, Bsep, and bile salt secretory failure in isolated rat hepatocyte couplets: a role for protein kinase C and prevention by protein kinase A. *Toxicol Sci.* 91, 150-158.

Pimplikar, S.W., and Simons, K. (1994) Activators of protein kinase A stimulate apical but not basolateral transport in epithelial Madin-Darby canine kidney cells. *J Biol Chem.* 269, 19054-19059.

Prekeris, R., Klumperman, J., and Scheller, R. H. (2000) A Rab11/Rip11 protein complex regulates apical membrane trafficking via recycling endosomes. *Mol Cell.* 6, 1437-1448.

Prince, L. S., Workman, R. B. Jr., and Marchase, R. B. (1994) Rapid endocytosis of the cystic fibrosis transmembrane conductance regulator chloride channel. *Proc Natl Acad Sci U S A.* 91, 5192-5196.

Prinz, A., Diskar, M., Erlbruch, A., Herberg, F. W. (2006) Novel, isotype-specific sensors for protein kinase A subunit interaction based on bioluminescence resonance energy transfer (BRET). *Cell Signal.* 18, 1616-1625.

Pullar, C. E., and Isseroff, R. R. (2006) The beta 2-adrenergic receptor activates pro-migratory and pro-proliferative pathways in dermal fibroblasts via divergent mechanisms. *J Cell Sci.* 119, 592-602.

Rabouille, C., Jokitalo, E. (2003) Golgi apparatus partitioning during cell division. *Mol Membr Biol.* 20, 117-127.

Rangarajan, S., Enserink, J. M., Kuiperij, H. B., de Rooij, J., Price, L. S., Schwede, F., and Bos, J. L. (2003) Cyclic AMP induces integrin-mediated cell adhesion through Epac and Rap1 upon stimulation of the beta 2-adrenergic receptor. *J Cell Biol.* 160, 487-493.

Razani, B., Rubin, C. S., and Lisanti, M. P. (1999) Regulation of cAMP-mediated signal transduction via interaction of caveolins with the catalytic subunit of protein kinase A. *J Biol Chem.* 274, 26353-26360.

Rizzolo, L. J., and Joshi, H. C. (1993) Apical orientation of the microtubule organizing center and associated gamma-tubulin during the polarization of the retinal pigment epithelium in vivo. *Dev Biol.* 157, 147-156.

Robin, P., Rossignol, B., and Raymond, M. N. (1998) PKA inhibitor, H-89, affects the intracellular transit of regulated secretory proteins in rat lacrimal glands. *Am J Physiol.* 274, 262-271.

Roelofsen, H., Soroka, C. J., Keppler, D., and Boyer, J. L. (1998). Cyclic AMP stimulates sorting of the canalicular organic anion transporter (Mrp2/cMoat) to the apical domain in hepatocyte couplets. *J Cell Sci.* 111, 1137-1145.

Roma, M. G., Milkiewicz, P., Elias, E., and Coleman, R. (2000) Control by signaling modulators of the sorting of canalicular transporters in rat hepatocyte couplets: role of the cytoskeleton. *Hepatology.* 32, 1342-1356.

Roma, M. G., Stone, V., Shaw, R., and Coleman, R. (1998) Vasopressin-induced disruption of actin cytoskeletal organization and canalicular function in isolated rat hepatocyte couplets: possible involvement of protein kinase C. *Hepatology.* 28, 1031-1041.

Sai, Y., Nies, A. T., and Arias, I. M. (1999) Bile acid secretion and direct targeting of mdr1-green fluorescent protein from Golgi to the canalicular membrane in polarized WIF-B cells. *J Cell Sci.* 112, 4535-4545.

Salas, P. J. I. (1999) Insoluble γ -tubulin-containing structures are anchored to the apical network of intermediate filaments in polarized CACO-2 epithelial cells. *J. Cell Biol.* 146, 645-657.

Schell, M. J. , Maurice, M. , Stieger, B., and Hubbard, A. L. (1992). 5'nucleotidase is sorted to the apical domain of hepatocytes via an indirect route. *J Cell Biol.* 119, 1173-1182.

Schmidt, P. H., Dransfield, D. T., Claudio, J. O., Hawley, R. G., Trotter, K. W., Milgram, S. L., and Goldenring, J. R. (1999) AKAP350, a multiply spliced protein kinase A-anchoring protein associated with centrosomes. *J Biol Chem.* 274, 3055-3066.

Seibert, F. S., Chang, X. B., Aleksandrov, A. A., Clarke, D. M., Hanrahan, J. W., and Riordan, J. R. (1999) Influence of phosphorylation by protein kinase A on CFTR at the cell surface and endoplasmic reticulum. *Biochim Biophys Acta.* 1461, 275-283.

Seino, S., and Shibasaki, T. (2005) PKA-dependent and PKA-independent pathways for cAMP-regulated exocytosis. *Physiol Rev.* 85, 1303-1342.

Shanks, R. A., Steadman, B. T., Schmidt, P. H., and Goldenring, J. R. (2002) AKAP350 at the Golgi apparatus. I. Identification of a distinct Golgi apparatus targeting motif in AKAP350. *J Biol Chem.* 277, 40967-40972.

Shen, X., Xu, K. F., Fan, Q., Pacheco-Rodriguez, G., Moss, J., and Vaughan, M. (2006) Association of brefeldin A-inhibited guanine nucleotide-exchange protein 2 (BIG2) with recycling endosomes during transferrin uptake. *Proc Natl Acad Sci U S A.* 103, 2635-2640.

Shi, G. X., Rehmann, H., Andres, D. A. (2006) A Novel cAMP-dependent Epac-Rit Signaling Pathway Contributes to PACAP38-mediated Neuronal Differentiation. *Mol Cell Biol.* [Epub ahead of print]

Shin, H. W., Morinaga, N., Noda, M., and Nakayama, K. (2004) BIG2, a guanine nucleotide exchange factor for ADP-ribosylation factors: its localization to recycling endosomes and implication in the endosome integrity. *Mol Biol Cell.* 15, 5283-5294.

Shorter, J., and Warren, G. (2002) Golgi architecture and inheritance. *Annu Rev Cell Dev Biol.* 18, 379-420.

Sietsma, H., Veldman, R. J., and Kok, J. W. (2001). The involvement of sphingolipids in multidrug resistance. *J Membr Biol.* 181, 153-162.

Singh, A. K., Tasken, K., Walker, W., Frizzell, R. A., Watkins, S. C., Bridges, R. J., and Bradbury, N. A. (1998) Characterization of PKA isoforms and kinase-dependent activation of chloride secretion in T84 cells. *Am J Physiol.* 275, 562-570.

Skálhegg, B. S. and Taskén, K. (1997) Specificity in the cAMP/PKA signaling pathway. Differential expression, regulation, and subcellular localization of subunits of PKA. *Frontiers in Bioscience.* 2, 331-342.

Skálhegg, B. S., Landmark, B., Foss, K. B., Lohmann, S. M., Hansson, V., Lea, T., and Jahnsen, T. (1992) Identification, purification, and characterization of subunits of cAMP-dependent protein kinase in human testis. Reverse mobilities of human RII alpha and RII beta on sodium dodecyl sulfate-polyacrylamide gel electrophoresis compared with rat and bovine RII α s. *J Biol Chem.* 267, 5374-5379.

Snyder, P.M. (2000). Liddle's syndrome mutations disrupt cAMP-mediated translocation of the epithelial Na(+) channel to the cell surface. *J Clin Invest.* 105, 45-53.

Sormunen, R., Eskelinen, S., Lehto, V. P. (1993) Bile canaliculus formation in cultured HEPG2 cells. *Lab Invest.* 68, 652-662.

Sudo, R., Mitaka, T., Ikeda, M., and Tanishita, K. (2005) Reconstruction of 3D stacked-up structures by rat small hepatocytes on microporous membranes. *FASEB J.* 19, 1695-1697.

Taouil, K., Hinnrasky, J., Hologne, C., Corlieu, P., Klossek, J. M., and Puchelle, E. (2003) Stimulation of beta 2-adrenergic receptor increases cystic fibrosis transmembrane conductance regulator expression in human airway epithelial cells through a cAMP/protein kinase A-independent pathway. *J Biol Chem.* 278, 17320-17327.

Taylor, S. S., Kim, C., Vigil, D., Haste, N. M., Yang, J., Wu, J., Anand, G. S. (2005) Dynamics of signaling by PKA. *Biochim Biophys Acta.* 1754, 25-37.

Thomas, C.P., Campbell, J.R., Wright, P.J., and Husted, R.F. (2004). cAMP-stimulated Na⁺ transport in H441 distal lung epithelial cells: role of PKA, phosphatidylinositol 3-kinase, and sgk1. *Am J Physiol Lung Cell Mol Physiol.* 287, 843-851.

Tietz, P. S., Marinelli, R. A., chen, X. M., Huang, B., Cohn, J., Kole, J., McNiven, M. A., Alper, S., and LaRusso, N. F. (2003). Agonist-induced coordinated trafficking of functionally related transport proteins for water and ions in cholangiocytes. *J Biol Chem.* 278, 20413-20419.

Togawa, A., Morinaga, N., Ogasawara, M., Moss, J., and Vaughan, M. (1999) Purification and cloning of a brefeldin A-inhibited guanine nucleotide-exchange protein for ADP-ribosylation factors. *J Biol Chem.* 274, 12308-12315.

Truschel, S. T., Wang, E., Ruiz, W. G., Leung, S. M., Rojas, R., Lavelle, J., Zeidel, M., Stoffer, D., and Apodaca, G. (2002). Stretch-regulated exocytosis/endocytosis in bladder umbrella cells. *Mol Biol Cell.* 13, 830-846.

Tuma, P. L., Nyasae, L. K., Backer, J. M., Hubbard, A. L. (2001) Vps34p differentially regulates endocytosis from the apical and basolateral domains in polarized hepatic cells. *J Cell Biol.* 154, 1197-1208.

Tuma, P. L. , Finnegan, C. M. , Yi, J. H. , and Hubbard, A. L. (1999). Evidence for apical endocytosis in polarized hepatic cells: phosphoinositide 3-kinase inhibitors lead to the lysosomal accumulation of resident apical plasma membrane proteins. *J Cell Biol.* 145, 1089-1102.

Valenti, G., Procino, G., Liebenhoff, U., Frigeri, A., Bendetti, P. A., Ahnert-Hilger, G., Nurnberg, B., Svelto, M., and Rosenthal, W. (1998). A

heterotrimeric G protein of the Gi family is required for cAMP-triggered trafficking of aquaporin 2 in kidney epithelial cells. *J Biol Chem.* 273, 22627-22634.

van Helvoort, A. , Smith, A. J. , Sprong, H. , Fritzsche, I. , Schinkel, A. H., Borst, P. , and van Meer, G. (1996). MDR1 P-glycoprotein is a lipid translocase of broad specificity, while MDR3 P-glycoprotein specifically translocates phosphatidylcholine. *Cell.* 87, 507-517.

van der Wouden, J. M., van IJzendoorn, S. C. D., and Hoekstra, D. (2002). Oncostatin M regulates membrane traffic and stimulates bile canalicular membrane biogenesis in HepG2 cells. *EMBO J.* 21, 6409-6418.

van IJzendoorn, S. C., Theard, D., Van Der Wouden, J. M., Visser, W., Wojtal, K. A., and Hoekstra, D. (2004) Oncostatin M-stimulated apical plasma membrane biogenesis requires p27(Kip1)-regulated cell cycle dynamics. *Mol Biol Cell.* 15, 4105-4114.

van IJzendoorn, S. C. D., van der Wouden, J. M., Liebisch, G., Schmitz, G., and Hoekstra, D. (2004). Polarized membrane traffic and cell polarity development is dependent on dihydroceramide synthase-regulated sphinganine turnover. *Mol Biol Cell.* 15, 4115-4124.

van IJzendoorn, S. C., and Hoekstra, D. (2000) Polarized sphingolipid transport from the subapical compartment changes during cell polarity development. *Mol.Biol.Cell* 11, 1093-1101.

van IJzendoorn, S. C., and Hoekstra, D. (1999) Polarized sphingolipid transport from the subapical compartment: evidence for distinct sphingolipid domains. *Mol Biol Cell.* 10, 3449-3461.

van IJzendoorn, S. C., and Hoekstra, D. (1998) (Glyco)sphingolipids are sorted in sub-apical compartments in HepG2 cells: a role for non-Golgi-related intracellular sites in the polarized distribution of (glyco)sphingolipids. *J Cell Biol.* 142, 683-696.

van IJzendoorn, S. C. D., and Hoekstra, D. (1998). (Glyco)sphingolipids are sorted in sub-apical compartments in HepG2 cells: a role for non-Golgi-related intracellular sites in the polarized distribution of (glyco)sphingolipids. *J Cell Biol.* 142, 683-396.

van IJzendoorn, S. C. D., Zeggars, M. M., Kok, J. W., and Hoekstra, D. (1997). Segregation of glucosylceramide and sphingomyelin occurs in the apical to basolateral transcytotic route in HepG2 cells. *J Cell Biol.* 137, 347-357.

Veit, B., Yucel, J. K., and Malhotra, V. (1993) Microtubule independent vesiculation of Golgi membranes and the reassembly of vesicles into Golgi stacks. *J Cell Biol.* 122, 1197-1206.

Vinciguerra, M., Hasler, U., Mordasini, D., Roussel, M., Capovilla, M., Ogier-Denis, E., Vandewalle, A., Martin, P. Y., and Feraille, E. (2005) Cytokines and sodium induce protein kinase A-dependent cell-surface Na,K-ATPase recruitment via dissociation of NF-kappaB/IkappaB/protein kinase A catalytic subunit complex in collecting duct principal cells. *J Am Soc Nephrol.* 16, 2576-2585.

Viollet, B., Kahn, A., and Raymondjean, M. (1997) Protein kinase A-dependent phosphorylation modulates DNA-binding activity of hepatocyte nuclear factor 4. *Mol Cell Biol.* 17, 4208-4219.

Viste, K., Kopperud, R. K., Christensen, A. E., and Doskeland, S. O. (2005) Substrate enhances the sensitivity of type I protein kinase a to cAMP. *J Biol Chem.* 280, 13279-13284.

Wald, F. A., Figueroa, Y., Oriolo, A. S., and Salas, P. J. (2003) Membrane repolarization is delayed in proximal tubules after ischemia-reperfusion: possible role of microtubule-organizing centers. *Am J Physiol Renal Physiol* 285, 230-240.

Wakabayashi, Y., Dutt, P., Lippincott-Schwartz, J., and Arias, I. M. (2005) Rab11a and myosin Vb are required for bile canaliculi formation in WIF-B9 cells. *Proc Natl Acad Sci U S A.* 102, 15087-15092.

Wang, W., Soroka, C. J., Mennone, A., Rahner, C., Harry, K., Pypaert, M., and Boyer, J. L. (2006) Radixin is required to maintain apical canalicular membrane structure and function in rat hepatocytes. *Gastroenterology.* 131, 878-884.

Wang EC, Lee JM, Ruiz WG, Balestreire EM, von Bodungen M, Barrick S, Cockayne DA, Birder LA, Apodaca G. (2005). ATP and purinergic receptor-dependent membrane traffic in bladder umbrella cells. *J Clin Invest.* 115, 2412-2422.

Wang, E., Brown, P. S., Aroeti, B., Chapin, S. J., Mostov, K. E., and Dunn, K. W. (2000) Apical and basolateral endocytic pathways of MDCK cells meet in acidic common endosomes distinct from a nearly-neutral apical recycling endosome. *Traffic.* 1, 480-493.

Ward, D. T., Hammond, T. G., and Harris, H. W. (1999). Modulation of vasopressin-elicited water transport by trafficking of aquaporin2-containing vesicles. *Annu Rev Physiol.* 61, 683-697.

Warren, G. (1995) Intracellular membrane morphology. *Philos Trans R Soc Lond B Biol Sci.* 349, 291-295.

Weinberger, A., Kamena, F., Kama, R., Spang, A., and Gerst, J. E. (2005) Control of Golgi morphology and function by Sed5 t-SNARE phosphorylation. *Mol Biol Cell.* 16, 4918-4930.

Westphal, R. S., Soderling, S. H., Alto, N. M., Langeberg, L. K., and Scott, J. D. (2000) Scar/WAVE-1, a Wiskott-Aldrich syndrome protein, assembles an actin-associated multi-kinase scaffold. *EMBO J.* 19, 4589-4600.

Wiley, J. C., Wailes, L. A., Idzerda, R. L., and McKnight, G. S. (1999) Role of regulatory subunits and protein kinase inhibitor (PKI) in determining nuclear localization and activity of the catalytic subunit of protein kinase A. *J. Biol Chem.* 274, 6381-6387.

Witczak, O., Skalhegg, B. S., Keryer, G., Bornens, M., Tasken, K., Jahnsen, T., and Orstavik, S. (1999) Cloning and characterization of a cDNA encoding an A-kinase anchoring protein located in the centrosome, AKAP450. *EMBO J.* 18, 1858-1868.

Wittmaack, F. M., Weber, W., and Hilz, H. (1983) Isoenzymes of cAMP-dependent protein kinase in developing rat liver and in malignant hepatic tissues. *Eur J Biochem.* 129, 669-674.

Wojtal, K. A., de Vries, E., Hoekstra, D., van IJzendoorn, S. C. (2006) Efficient trafficking of MDR1/P-glycoprotein to apical canalicular plasma membranes in HepG2 cells requires PKA-Rilalpha anchoring and glucosylceramide. *Mol Biol Cell.* 17, 3638-3650.

Wong, W., and Scott, J. D. (2004) AKAP signalling complexes: focal points in space and time. *Nat Rev Mol Cell Biol.* 5, 959-970.

Yang, S., Fransson, U., Fagerhus, L., Holst, L. S., Hohmeier, H. E., Renstrom, E., and Mulder, H. (2004) Enhanced cAMP protein kinase A signaling determines improved insulin secretion in a clonal insulin-producing beta-cell line (INS-1 832/13). *Mol Endocrinol.* 18, 2312-2320.

Yao, X., and Forte, J.G. (2003). Cell biology of acid secretion by the parietal cell. *Annu Rev Physiol.* 65, 103-131.

Zaal, K. J. , Kok, J. W. , Sormunen, R. , Eskelinen, S. , and Hoekstra, D. (1994). Intracellular sites involved in the biogenesis of bile canaliculi in hepatic cells. *Eur J Cell Biol.* 63, 10-19.

Zegers, M. M., and Hoekstra, D. (1998) Mechanisms and functional features of polarized membrane traffic in epithelial and hepatic cells. *Biochem J.* 336, 257-269.

Zegers, M. M., Zaal, K. J., van IJzendoorn, S. C., Klappe, K., and Hoekstra, D. (1998) Actin filaments and microtubules are involved in different membrane traffic pathways that transport sphingolipids to the apical surface of polarized HepG2 cells. *Mol Biol Cell.* 9, 1939-1949.

Zegers, M.M. , Zaal, K.J. , and Hoekstra, D. (1998). Functional involvement of proteins, interacting with sphingolipids, in sphingolipid transport to the

canalicular membrane in the human hepatocytic cell line, HepG2? *Hepatology*. 27, 1089-1097.

Zegers, M.M., and Hoekstra, D. (1997). Sphingolipid transport to the apical plasma membrane domain in human hepatoma cells is controlled by PKC and PKA activity: a correlation with cell polarity in HepG2 cells. *J Cell Biol.* 138, 307-321.

Zegers, M. M., Kok, J. W., and Hoekstra, D. (1997) Use of photoactivatable sphingolipid analogues to monitor lipid transport in mammalian cells. *Biochem J.* 328, 489-498.

Zeligs, J. D. (1979) Association of centrioles with clusters of apical vesicles in mitotic thyroid epithelial cells. Are centrioles involved in directing secretion? *Cell Tissue Res.* 201, 11-21.

Zhou, Y. P., Marlen, K., Palma, J. F., Schweithzer, A., Reilly, L., Gregoire, F. M., Xu, G. G., Blume, J. E., and Johnson, J. D. (2003). Overexpression of repressive cAMP response element modulators in high glucose and fatty acid-treated rat islets. A common mechanism for glucose toxicity and lipotoxicity? *J Biol Chem.* 278, 51316-51323.